

Robustness represents the ability of the surveillance system to produce acceptable outcomes over a range of assumptions about uncertainty by maximizing the reliability of an adequate outcome.

	Surveillance design step	Advice for improvement of ROBUSTNESS
1	Surveillance system	
1.1	Hazard	
1.2	Surv. Objective	
1.3	Geographical area covered	
1.4	Susceptible species	
1.5	Risk characteristics	
2	Components overview	
3	Target population	
3.1	Target species	
3.2	Target sector	
3.3	Sectors missed	
3.4	Geographical area covered	
3.5	Target criteria	
3.6	Percentage covered	
4	Disease suspicion	
4.1	Definition	
4.2	Obligations	
4.3	Notification procedures	
4.4	Actions upon suspicions	
4.5	Actions upon confirmation	
5	Enhancements	
6	Testing protocol	Consider choosing a test with low uncertainty in sensitivity and specificity to increase robustness.
6.1	Type of test to be carried out	
6.2	Type of sample to be collected	
6.3	Pooling	
6.4	Screening/first test	
6.5	Confirmatory/ second test	
6.6	Further details	

7	Study design	If you are uncertain about input values (e.g. for sample size, test characteristics) or assumptions made about the population, run a sensitivity analysis to determine the effectiveness of your components over the full range of values. If any parameter has a major effect on the outcome, consider collecting additional data to minimize uncertainty for this parameter.
7.1	Point of sample collection	
7.2	Selection of units	
7.3	Target unit	
7.4	Sampling unit	
7.5	Sampling design	
7.6	Number of units in the target population	
7.7	Sensitivity of the testing protocol	
7.8	Specificity of the testing protocol	
8	Sampling strategy	
8.1	Sampling at the primary sampling unit (PSU) level:	
8.2	Sampling at the secondary sampling unit (SSU) level:	
8.3	Selection criteria WITHIN the population	
8.4	Risk-based allocation	
8.5	Sample size calculation	
8.6	Sample allocation at the primary level	
8.7	Sample allocation at the Secondary level	
8.8	Sample collection timeline	
9	Data Generation/ Sampling collection process	
9.1	WHO will collect the samples?	
9.2	HOW will samples be collected?	
9.3	WHEN/HOW OFTEN will samples be collected?	
9.4	Training	
9.5	Follow-up	
10	Transfer means	
10.1	HOW will samples be transferred?	If complicated procedures for transportation are required, robustness may be reduced.
10.2	WHEN/HOW OFTEN will samples be collected?	
10.3	Training	
11	Data Translation/ sample analyses process	
11.1	WHO will perform the analyses?	
11.2	HOW will samples be analysed	
11.3	WHEN/HOW OFTEN will samples be collected?	

11.4	Expected LOAD	
11.5	Training	
11.6	Follow-up	
12	Epidemiological analyses	
12.1	Are there any epidemiological DATA that need to be collected?	
12.2	WHO will perform the analyses?	
12.3	HOW will epidemiological analyses be performed?	
12.4	WHEN/HOW OFTEN?	
12.5	Training	
12.6	Data management needs	
12.7	Software needs	
13	Dissemination of results	
13.1	WHO will disseminate the results?	
13.2	WHO is the TARGET of dissemination?	
13.3	HOW will results be disseminated?	
13.4	WHEN/HOW OFTEN?	
14	Surveillance review	
14.1	Who	
14.2	When	
14.3	How often	



From:

<https://survtools.org/wiki/surveillance-design-framework/> - **Surveillance Design Framework Wiki**

Permanent link:

<https://survtools.org/wiki/surveillance-design-framework/doku.php?id=robustness&rev=1443782480>

Last update: **2015/10/02 12:41**

